


CBCS SCHEME



USN

18MAT31

Third Semester B.E. Degree Examination, July/August 2021 Transform Calculus, Fourier Series and Numerical Techniques

Time: 3 hrs.

Max. Marks:100

Note: Answer any FIVE full questions.

- 1 a. Find $L[t e^{-2t} \sin 4t]$. (06 Marks)
 - b. A periodic function of period $2\pi/\omega$ is defined by $f(t) = \begin{cases} E \sin \omega t, & 0 \leq t < \pi/\omega \\ 0, & \pi/\omega \leq t < 2\pi/\omega \end{cases}$. Where E and ω are constants. (07 Marks)
 - c. Solve : $y''(t) + k^2 y(t) = 0$; $y(0) = 0$ and $y'(0) = 1$ by Laplace transformation. (07 Marks)
 - 2 a. Find : i) $L^{-1}\left\{\frac{s^2 - 3s + 4}{s^3}\right\}$ ii) $L^{-1}\left[\text{Cot}^{-1}\left(\frac{S}{2}\right)\right]$. (06 Marks)
 - b. Find the inverse Laplace transform of $\frac{1}{(s-1)(s^2+1)}$ by using convolution theorem. (07 Marks)
 - c. Express the following function in terms of Heaviside step function and hence find its Laplace transform where $f(t) = \begin{cases} t^2, & 0 < t \leq 2 \\ 4t, & t > 2 \end{cases}$. (07 Marks)
 - 3 a. Expand $f(x) = x(2\pi - x)$ as a Fourier series in $[0, 2\pi]$. (06 Marks)
 - b. Obtain Fourier series for the function $f(x)$ given by $f(x) = \begin{cases} 1 + \frac{2x}{\pi}, & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi}, & 0 \leq x \leq \pi \end{cases}$. (07 Marks)
 - c. Find the half range sine series of $f(x) = \frac{e^{ax}}{\sinh a\pi}$ in $(0, \pi)$. (07 Marks)
 - 4 a. Find the Fourier series expansion of $f(x)$ given by $f(x) = \begin{cases} 1 & 0 < x < 1 \\ 2 & 1 < x < 3 \end{cases}$. (06 Marks)
 - b. Find the half range sine series for x^2 in $(0, \pi)$. (07 Marks)
 - c. The values of x and the corresponding values of $f(x)$ over a period T are given below. Show that $f(x) = 0.75 + 0.37 \cos \theta + 1.004 \sin \theta$ where $\theta = \frac{2\pi x}{T}$. (07 Marks)
- | | | | | | | | |
|------|------|------|------|------|-------|-------|------|
| x | 0 | T/6 | T/3 | T/2 | 2T/3 | 5T/6 | T |
| f(x) | 1.98 | 1.30 | 1.05 | 1.30 | -0.88 | -0.25 | 1.98 |
- 5 a. State: i) Initial and final value theorems ii) Find the Z -transform of $\cos\left(\frac{n\pi}{2} + \frac{\pi}{4}\right)$. (06 Marks)
 - b. Find the complex Fourier transform of the function $f(x) = \begin{cases} 1 & \text{for } |x| \leq a \\ 0 & \text{for } |x| > a \end{cases}$.
Hence evaluate $\int_0^\infty \left(\frac{\sin x}{x}\right) dx$. (07 Marks)
 - c. Compute the inverse Z-transform of $\frac{3z^2 + 2z}{(5z-1)(5z+2)}$. (07 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.
2. Any revealing of identification, appeal to evaluator and /or equations written eg. 42+8=50, will be treated as malpractice.

- 6 a. Find the Fourier cosine transform of $f(x) = \begin{cases} x, & 0 < x < 2 \\ 0, & \text{else where} \end{cases}$. (06 Marks)
- b. Find the Z-transform of $2n + \sin \frac{n\pi}{4} + 1$. (07 Marks)
- c. Solve the difference equation : $u_{n+2} - 3u_{n+1} + 2u_n = 0$, with $u_0 = 0$ and $u_1 = -1$. (07 Marks)
- 7 a. Find by Taylor's series method the value of y at $x = 0.1$ to five places of decimals from $\frac{dy}{dx} = x^2y - 1, y(0) = 1$. (06 Marks)
- b. Use fourth order Runge-Kutta method to solve $(x + y)\frac{dy}{dx} = 1, y(0.4) = 1$ at $x = 0.5$ correct to four decimal places. (07 Marks)
- c. If $\frac{dy}{dx} = 2e^x - y, y(0) = 2, y(0.1) = 2.010, y(0.2) = 2.040$ and $y(0.3) = 2.090$, find $y(0.4)$ correct to four decimal places by using Milne's predictor – corrector method and applying corrector formula twice. (07 Marks)
- 8 a. Using modified Euler's formula compute $y(1.1)$ correct to five decimal places given that $\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}$ and $y = 1$ at $x = 1$. [taking $h = 0.1$]. (06 Marks)
- b. Employ Taylor's series method to find y at $x = 0.1$ and 0.2 correct to four places of decimal. Given $\frac{dy}{dx} - 2y = 3e^x, y(0) = 0$. (07 Marks)
- c. Solve the differential equation $y' + y + xy^2 = 0$ with the initial values of $y : y_0 = 1, y_1 = 0.9008, y_2 = 0.8066, y_3 = 0.722$ corresponding to the values of $x : x_0 = 0, x_1 = 0.1, x_2 = 0.2, x_3 = 0.3$ by computing the value of y corresponding to $x = 0.4$ applying Adams – Bashforth predictor and corrector formula. (07 Marks)
- 9 a. Given $y'' - xy' - y = 0$ with the initial conditions $y(0) = 1, y'(0) = 0$, compute $y(0.2)$ using fourth order Runge-Kutta method. (06 Marks)
- b. Derive Euler's equation in the standard form $\frac{\partial f}{\partial y} - \frac{d}{dx} \left(\frac{\partial f}{\partial y'} \right) = 0$. (07 Marks)
- c. A heavy cable hangs freely under gravity between two fixed points. Show that the shape of the cable is a catenary. (07 Marks)
- 10 a. Apply Milne's method to compute $y(0.8)$ given that $y'' = 1 - 2yy'$ and the following table of initial values. (07 Marks)
- | | | | | |
|----|---|--------|--------|--------|
| x | 0 | 0.2 | 0.4 | 0.6 |
| y | 0 | 0.02 | 0.0795 | 0.1762 |
| y' | 0 | 0.1996 | 0.3937 | 0.5689 |
- b. Prove that the geodesics on a plane are straight line. (06 Marks)
- c. Find the extremal of the functional : $\int_{x_0}^{x_1} (y^2 + y'^2 - 2y \sin x) dx$. (07 Marks)
